



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : C12N 15/10, 15/62, 15/36, 15/51, 7/00, C07K 7/08, A61K 39/12, G01N 33/577		A2	(11) International Publication Number: WO 94/26886 (43) International Publication Date: 24 November 1994 (24.11.94)
(21) International Application Number: PCT/IT94/00054 (22) International Filing Date: 5 May 1994 (05.05.94) (30) Priority Data: RM93A000301 11 May 1993 (11.05.93) IT (71) Applicant (for all designated States except US): ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P. ANGELETTI S.P.A. [IT/IT]; Via Pontina Km 30.600, I-00040 Pomezia (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): FELICI, Franco [IT/IT]; Viale Bruno Rizzieri, 247, I-00173 Roma (IT). LUZZAGO, Alessandra [IT/IT]; Via Beata Vergine del Carmelo, 54, I-00144 Roma (IT). NICOSIA, Alfredo [IT/IT]; Via Beata Vergine del Carmelo, 54, I-00144 Roma (IT). MONACI, Paolo [IT/IT]; Via Appia Nuova, 420, I-00181 Roma (IT). CORTESE, Riccardo [IT/IT]; Via Massimiliano Massimo, 16, I-00144 Roma (IT). (74) Agents: DI CERBO, Mario et al.; Società Italiana Brevetti S.p.A., Piazza di Pietra, 39, I-00186 Roma (IT).		(81) Designated States: AU, BR, CA, CN, JP, RU, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BI, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  Published Without international search report and to be republished upon receipt of that report.	
(54) Title: PROCESS FOR THE PREPARATION OF IMMUNOGENS OR DIAGNOSTIC REAGENTS, AND IMMUNOGENS OR DIAGNOSTIC REAGENTS THEREBY OBTAINABLE			
<p>lac promoter      leader sequence      pVIII      α-peptide</p> <p>GAATTC.TAG.GATCCC GluPhe.END.AspPro EcoRI      BamIII</p>			
(57) Abstract			
<p>A process for the preparation of immunogens or diagnostic reagents that mimic an antigen or a pathogenic organism specific to a disease, essentially characterized by the following operations: identification of at least one antibody that reacts with the antigen or pathogenic organism specific to the disease; construction of phage libraries which display on the surface of the capsid oligopeptides, expressed from random sequence oligonucleotide inserts introduced into a gene coding for a phage capsid protein using genetic manipulation techniques (for example using a plasmid engineered for the purposes of the invention, the genetic map of which is shown in the figure); selection of the phages that display on the surfaces of the capsid antigenic oligopeptides recognized by said antibody; optional use of the selected phages and/or fragments thereof and/or their derivatives for the formulation of diagnostic kits for the specific pathogenic agent, or in general for the diseases, including immunological disorders typical of so-called autoimmune diseases, with known or unknown etiology and/or pathogenesis; optional use of the selected phages and/or fragments thereof and/or their derivatives for the formulation of an antagonist of the antigen-antibody reactions for treatment of the disease induced by said antigen; optional use of the selected phages and/or fragments thereof and/or their derivatives to induce a tolerance of the phenomena of hypersensitivity and/or allergy to compounds and/or natural or synthetic preparations; optional immunization of an organism by means of the selected phages and/or fragments thereof and/or their derivatives; and optional verification of the presence, in the serum of the immunized organism, of antibodies that recognize the above antigen or organism specific to the disease.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MD	Republic of Moldova	TJ	Tajikistan
DE	Germany	MG	Madagascar	TT	Trinidad and Tobago
DK	Denmark	ML	Mali	UA	Ukraine
ES	Spain	MN	Mongolia	US	United States of America
FI	Finland			UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

PROCESS FOR THE PREPARATION OF IMMUNOGENS OR DIAGNOSTIC REAGENTS, AND IMMUNOGENS OR DIAGNOSTIC REAGENTS THEREBY OBTAINABLE

DESCRIPTION

The subject of the present invention is a process for the preparation of immunogens or diagnostic reagents that mimic an antigen or a pathogenic organism specific to a disease, even if this is uncharacterized or even unknown (thereby including auto-immune diseases whose etiology and/or pathogenesis is known or unknown). This process is based on the existence and availability of antibodies, both monoclonal or polyclonal, or of serum containing antibodies, which react specifically with the organism causing the infection.

Antibodies suitable for use in this process can be specific for any antigen of interest for which an immunogen or diagnostic reagent that mimics the antigen is sought. The antigen can be a protein or peptide whether synthetic, derived from a natural source, or produced recombinantly; carbohydrate; polysaccharide; glycoprotein; hormone; receptor; antibody; virus; substrate; metabolite; transition state analog; cofactor; drug; dye; nutrient; growth factor; cellular component; oncogene product; bacteria and their extracellular products; mammalian cells and extracts therefrom including tumor cells, virus infected cells and normal cells; parasites; protozoa; malarial antigens; helminths; fungi; rickettsia; or an allergen including but not limited to pollens, dusts, danders or extracts of the same; or a venom, poison, toxin, or toxoid; nucleic acids including DNA; or any other antigen without limitation. Antigens of viruses which are suitable for use in the present invention include antigens from the viruses including but not limited to polio virus, influenza virus, HIV, HTLV, papilloma virus, adeno virus, parainfluenza virus, measles virus,

- 2 -

mumps virus, respiratory syncytial virus, shipping fever virus, Western and Eastern encephalomyelitis virus, Japanese B encephalomyelitis virus, Russian spring-summer encephalomyelitis virus, hog cholera virus, hepatitis virus, pox virus, rabies, virus, distemper virus, herpes virus, cytomegalo virus, foot and mouth disease virus, rhinovirus, Newcastle disease virus, vaccinia virus; and pseudorabies virus. The mime can be an immunogen, a vaccine, an inhibitor or activator, etc. without limitation.

As is known, all vaccines and diagnostic reagents currently on sale or undergoing clinical tests are conventionally obtained by means of processes based on the manipulation, modification and/or adaptation of pathogenic organisms or components thereof. These methods have given good results, but are not without problems. The greatest limitation associated with these methods is connected with the fact that they depend upon the availability of information and/or material directly deriving from pathogenic organisms or components thereof.

Previous attempts to overcome the above described limitation have so far failed for a lack of an efficient and reproducible experimental protocol; in particular, these attempts did not provide sufficient information in order to identify and characterize immunogenic mimics to be used for diagnosis and vaccine therapy. It must be emphasized that the present invention is focused on the development of a new technology aimed at overcoming conceptual and technical inadequancies of previously proposed protocols.

A key feature of the present invention is a novel strategy for the selection of antigenic and immunogenic mimics, based on the use, as reagents, of serum samples from patients and a counter-selection step utilizing serum samples from healthy individuals.

Use of the process for preparation according to

- 3 -

the present invention allows this limitation to be overcome, furthermore offering additional advantages which will be clear from the following description.

The process for the preparation of immunogens or diagnostic reagents that mimic an antigen or a pathogenic organism specific to a disease - according to the present invention - is essentially characterized by the following operations:

- identification of at least one antibody that reacts with the antigen or pathogenic organism specific to the disease;

- construction of phage libraries which display on the surface of the capsid oligopeptides, expressed from random sequence oligonucleotidic inserts introduced into a gene coding for the phage capsid using genetic manipulation techniques;

- selection of the phages that display on the surfaces of the capsid antigenic oligopeptides recognized by said antibody; optionally by selection with a first pathologic serum, subsequent screening with a second different pathologic serum and counter-screening with a panel of sera from healthy individuals;

- optional use of the selected phages and/or fragments thereof and/or their derivatives for the formulation of diagnostic kits for the specific pathogenic agent, or in general for the diseases, including immunological disorders typical of so-called autoimmune diseases, with known or unknown etiology and/or pathogenesis;

- optional use of the selected phages and/or fragments thereof and/or their derivatives for the formulation of an antagonist of the antigen-antibody reactions for treatment of the disease induced by said antigen;

- optional use of the selected phages and/or fragments thereof and/or their derivatives to induce a

- 4 -

tolerance of the phenomena of hypersensitivity and/or allergy to compounds and/or natural or synthetic preparations;

- optional immunization of an organism by means of the selected phages and/or fragments thereof and/or their derivatives; and

- optional verification of the presence, in the serum of the immunized organism, of antibodies that recognize the above antigen or organism specific to the disease.

Should said antibodies specific to said pathogenic agent also be protective or neutralizing, the material used for said immunization can be used to formulate a vaccine against said specific pathogenic agent.

The construction of phage libraries, according to the present invention, can be advantageously performed using the filamentous phages M13, F1 and Fd, or derivatives thereof. The reasons for this are the following:

- filamentous phages are commonly used as molecular vectors in the field of molecular biology and genetic engineering. For example, by taking advantage of their feature to contain a genome with a single DNA helix, they have been particularly used in DNA sequencing experiments, in direct site mutagenesis experiments and for the expression of proteins and peptides;

- the information required and sufficient for encapsidation of a single chain DNA genome has been well characterized and can be transferred to other molecular vectors;

- it has likewise been demonstrated that at least two proteins of the capsid of filamentous phages can be modified by means of the addition or insertion of additional amino acid sequences. The resulting phages are encapsidated, maintain their ability to replicate and, in most cases, to infect bacterial cells. The

- 5 -

foreign amino acid sequences are displayed on the surface of the phage, and can be recognized by interaction with antibodies or with other specific molecules according to the case.

The antibodies that can be used in the process for the preparation of immunogens and diagnostic reagents according to the present invention can be monoclonal antibodies, polyclonal antibodies, or antibodies contained in sera. The latter form of embodiment is of particular interest, because it provides for the first time a reproducible experimental strategy to identify novel antigenic and immunogenic mimics in absence of any information on the structure and properties of the natural and pathological antigen.

The gene coding for the phage capsid, with random sequence oligonucleotidic inserts, can be the gene coding for the protein VIII of the phage capsid or the gene coding for the protein III of said capsid.

The process according to the invention can be applied without restriction to any antibody or organism responsible for illness. Good results have been obtained using monoclonal antibodies, or sera specific for the surface antigen of the human hepatitis B virus (HBsAg).

The antigenic oligopeptides recognized by the antibodies used can be obtained by expression from random sequence oligonucleotidic inserts, using as a vector, for example, the plasmid pC89.

In the process for the preparation of immunogens and diagnostic reagents according to the present invention, it is possible to select phages containing in their capsid from the site identifying the restriction enzyme EcoRI (GAATTC) to that identifying the restriction enzyme BamHI (GGATCC), one of the aminoacidic sequences SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 7 and SEQ ID NO: 9 to 68.

The present invention is not limited to the

- 6 -

process for the preparation of immunogens or diagnostic reagents against a specific pathogenic agent, but also extends to the immunogens and diagnostic reagents obtainable using the process illustrated above, and to the phages usable in the process mentioned above.

Furthermore, the invention also extends to the plasmids pC89 containing, wholly or in part, a nucleotidic sequence chosen from the group comprising the sequences SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6 and SEQ ID NO:8.

The immunogens obtained from the process of the present invention are useful as vaccines or immunizing agents as well as being useful as diagnostic reagents. The vaccines or immunizing agents are administered to a patient in need of such treatment according to standard methods known in the art. The vaccine or immunizing agents can be administered and used either singly or in combination. The vaccines and immunogens of the present invention can comprise the phage or protein and peptides isolated therefrom.

Kits containing the immunogens obtained from the process of the present invention may be prepared. Such kits are used to detect the presence of the antigen in a sample. Such characterization is useful for a variety of purposes including but not limited to forensic analyses and epidemiological studies. Such a kit would comprise a compartmentalized carrier suitable to hold in close confinement at least one container. The carrier would further comprise reagents such as the immunogens, and antibodies suitable for detecting the antigens. The carrier may also contain a means for detection such as labeled antigen or enzyme substrates or the like.

Pharmaceutically useful compositions comprising the immunogens and vaccines of the present invention, may be formulated according to known methods such as by the admixture of a pharmaceutically acceptable carrier.



- 7 -

Examples of such carriers and methods of formulation may be found in Remington's Pharmaceutical Sciences. To form a pharmaceutically acceptable composition suitable for effective administration, such compositions will contain an effective amount of the vaccine or immunogen of the present invention.

Therapeutic or diagnostic compositions of the invention are administered to an individual in amounts sufficient to treat or diagnose the relevant disorders. The effective amount may vary according to a variety of factors such as the individual's condition, weight, sex and age. Other factors include the mode of administration. The pharmaceutical compositions may be provided to the individual by a variety of routes such as subcutaneous, topical, oral and intramuscular.

The present invention also has the objective of providing suitable topical, oral systemic and parenteral pharmaceutical formulations for use in the novel methods of treatment of the present invention. The compositions containing vaccine or immunogens identified according to this invention as the active ingredient can be administered in a wide variety of therapeutic dosage forms in conventional vehicles for administration. For example, the vaccines and immunogens can be administered in such oral dosage forms as tablets, capsules (each including timed release and sustained release formulations), pills, powders, granules, elixirs, tinctures, solutions, suspensions, syrups and emulsions, or by injection. Likewise, they may also be administered in intravenous (both bolus and infusion), intraperitoneal, subcutaneous, topical with or without occlusion, or intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts. An effective but non-toxic amount of the vaccines and immunogens can be employed.

The daily dosage of the vaccines and immunogens

- 8 -

may be varied over a wide range from 0,01 to 1000 mg per adult human/per day. For oral administration, the vaccines and immunogens are preferably provided in the form of scored or unscored tablets containin 0,01, 0,05, 0,1, 0,5, 1,0, 2,5, 5,0, 10,0, 15,0, 25,0, and 50,0 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the vaccines and immunogens is ordinarily supplied at a dosage level of from about 0,0001 mg/kg to about 100 mg/kg of body weight per day. The range is more particularly from about 0,001 mg/kg to 10 mg/kg of body weight per day. The dosages of the vaccines and immunogens are adjusted when combined to achieve desired effects. On the other hand, dosages of these various agents may be independently optimized and combined to achieve a synergistic result wherein the pathology is reduced more than it would be if either agent were used alone.

Advantageously, vaccine or immunogens of the present invention may be administered in a single dose, or a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, vaccine or immunogens for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

For combination treatment with more than one active agent, where the active agents are in separate dosage formulations, the active agents can be administered concurrently, or they each can be administered at separately staggered times.

The dosage regimen utilizing the vaccine or

- 9 -

immunogens of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular vaccine or immunogen thereof employed. A physician or veterinarian of ordinary skill can readily determine and prescribe the effective amount of the vaccine or immunogens of the present invention required to prevent, counter or arrest the progress of the condition. Optimal precision in achieving concentrations of the vaccine or immunogens of the present invention within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the vaccine or immunogens of the present invention availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of the vaccine or immunogens of the present invention.

In the methods of the present invention, the vaccine or immunogens herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active vaccine or immunogen component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders

- 10 -

include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include, without limitation, sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methylcellulose, agar bentonite, xanthan gum and the like.

For liquid forms the active vaccine or immunogen component can be combined in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methylcellulose and the like. Other dispersing agents which may be employed include glycerin and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

Topical preparations containing the active vaccine or immunogen component can be admixed with a variety of carrier materials well known in the art, such as, e.g. alcohols, aloe vera gel, allantoin, glycerine, vitamin A and E oils, mineral oil, PPG2 myristyl propionate, and the like to form, e.g. alcoholic solutions, topical cleansers, cleansing creams, skin gels, skin lotions, and shampoos in cream or gel formulations.

The vaccine or immunogens of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Vaccine or immunogens of the present invention may

- 11 -

also be delivered by the use of monoclonal antibodies as individual carriers to which the vaccine or immunogen molecules are coupled. The vaccine or immunogens of the present invention may also be coupled with soluble polymers as targetable vaccine or immunogen carriers. Such polymers can include polyvinyl-pyrrolidone, pyran copolymer, polyhydroxypropylmethacryl-amidephenol, polyhydroxyethylaspartamidephenol, or polyethyl-eneoxidepolylysine substituted with palmitoyl residues. Furthermore, the vaccine or immunogens of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a vaccine or immunogen, for example, polyactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydro-pyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

Up to this point a general description has been given of the subjects of the present invention. With the assistance of the following examples, a more detailed description will now be given of specific embodiments of the invention, aimed at giving a better understanding of the objects, characteristics, advantages and methods of application thereof. The following examples refer, respectively, to embodiments of the process for the preparation of immunogens and diagnostic reagents against a specific pathogenic agent, according to the present invention; to a demonstration of the effectiveness of the clones selected for preparation of immunogens based on the effects produced by samples of serum from test animals immunized using the single clones; and to a demonstration of the effectiveness of the clones selected for preparation of diagnostic reagents, based on their specific reaction with the serum of individuals immunized with HBsAg.

The single figure enclosed shows a portion of the

- 12 -

genetic map of the plasmid pC89 engineered for the purposes of the invention. The nucleotidic sequence for the restriction sites and the corresponding aminoacidic sequence are illustrated below the portion of the above mentioned genetic map. The wild-type aminoacidic sequence of the amino-terminal end of mature pVIII has been modified in order to introduce single EcoRI and BamHI sites.

#### Example 1

#### Process for the preparation of specific diagnostic reagents and immunogens for the disease caused by the human Hepatitis B virus (HBV)

The following process was used according to the invention:

A first "library" of epitopes was prepared, made up of phages having oligopeptides with 9 amino acids displayed on the surface of the capsid, expressed by random sequence oligonucleotidic inserts introduced into the gene coding for the protein VIII of the phage capsid. For expression of the recombinant proteins containing the epitope, the plasmid pC89 was used, engineered as described in F. Felici et al., Selection of Antibody Ligands from a Large Library of Oligopeptides Expressed on a Multivalent Exposition Vector, J. Mol. Biol. (1991), 222, 301-310, of which a portion of the genetic map is shown in the figure.

A second epitope "library" was prepared in the same way as the first, with the difference that two cysteine residues are present at the two ends of the insert, as described in A. Luzzago et al., Mimicking of discontinuous epitopes by phage displayed peptides, I. Epitope mapping of human H Ferritin using a phage library of constrained peptides, Gene (1993), 128, 51-57.

A sample of sera from individuals immunized using a recombinant form of the HBV surface antigen (HBsAg) was tested for the presence of specific antibodies

- 13 -

against HBsAg. The sera with a high antibody content against HBsAg were then used.

The antibodies contained in these sera were immobilized on a solid matrix and incubated with the phage libraries. The phages specifically retained by these antibodies were then eluted and amplified.

Bacteria infected by the phages selected as above were plated on a solid culture medium and transferred onto nitrocellulose filters. Subsequently, these filters were incubated with sera from individuals immunized against HBsAg different from the serum used for the first enrichment. The phages specifically recognized by antibodies present in these sera were identified and isolated.

The phages identified as above were counter-selected for reactivity to antibodies present in the sera of individuals not immune to HBsAg, using the ELISA test.

The phages resulting from this counter-selection were checked for specificity of reaction to anti-HBsAg antibodies by means of competition in the ELISA test.

The nucleotidic sequence coding for the epitopes identified in this manner was subsequently determined (SEQ ID NO: 4, 5, 6, 8).

#### EXAMPLE 2

Demonstration of the effectiveness of the selected phages as specific immunogens for the disease caused by the human hepatitis virus (HBV)

The following process was used:

The phages, selected as described above, were amplified and purified.

The purified phages were injected into test animals (mice, rats and rabbits), according to the process described by V.F. de la Cruz et al., Immunogenicity and Epitope Mapping of Foreign Sequences via Genetically Engineered Filamentous Phage, J. Biol. Chem. (1988) 263, 9, 4318-4322, and by J. Greenwood et

- 14 -

al., Multiple Display of Foreign Peptides on a Filamentous Bacteriophage, J. Mol. Biol. (1991) 220, 821-827.

The sera of animals immunized as above were tested for the presence of antibodies capable of interacting with HBsAg, using the ELISA test. This method underlined the presence of a high level of anti-HBsAg antibodies in the serum of test animals immunized as above.

The same immunization process was adopted, using as immunogens synthetic oligopeptides reproducing the amino acid sequence (SEQ ID from NO: 1 to NO: 3 and NO:7) of the epitopes identified according to the procedure given above. This method underlined the presence of a high level of anti-HBsAg antibodies in the serum of test animals immunized as above.

The same immunization process was adopted, using as immunogens recombinant forms produced in bacteria of the heavy chain of human ferritin displaying on their surface the synthetic oligopeptides reproducing the amino acid sequence of the epitopes identified according to the procedure given above. This method underlined the presence of a high level of anti-HBsAg antibodies in the serum of test animals immunized as above.

### EXAMPLE 3

Demonstration of the effectiveness of the selected phages as diagnostic reagents for determination of the presence of anti-HBsAg antibodies in serum.

A process comprising the following operations was used:

20 human sera from individuals vaccinated against HBsAg were analyzed in an ELISA test for their ability to recognize specifically the phage clones, identified as described in example 1, showing the peptide sequences described (SEQ ID from NO: 1 to NO: 3 and NO: 7). The results show that 80% of the sera



- 15 -

specifically recognize at least one of the four sequences selected.

16 human serums from patients suffering from hepatitis B and containing anti-HBsAg antibodies were analyzed in an ELISA test for their ability to recognize specifically the phage clones, identified as described in example 1, showing the peptide sequences described (SEQ ID from NO: 1 to NO: 3 and NO:7). The results show that 44% of the sera specifically recognize at least one of the four sequences selected.

20 human sera from individuals not vaccinated against HBsAg were analyzed in an ELISA test for their ability to recognize specifically the phage clones, identified as described in example 1, displaying the peptide sequences described (SEQ ID from NO: 1 to NO:3 and NO:7). The results show that none of the sera specifically recognize any of the four sequences selected.

#### Example 4

#### Process for the preparation of specific diagnostic reagents and immunogens for the disease caused by the human Hepatitis C virus (HCV)

The following process was used according to the invention:

A first "library" of epitopes was prepared, made up of phages having oligopeptides with 9 amino acids displayed on the surface of the capsid, expressed by random sequence oligonucleotidic inserts introduced into the gene coding for the protein VIII of the phage capsid. For the expression of the recombinant proteins containing the epitope, the plasmid pC89 was used, engineered as described in F. Felici et al., Selection of Antibody Ligands from a Large Library of Oligopeptides Expressed on a Multivalent Exposition Vector, J. Mol. Biol. (1991), 222, 301-310, of which a portion of the genetic map is shown in the figure.

A second epitope "library" was prepared in the

- 16 -

same way as the first, with the difference that two cysteine residues are present at the two ends of the insert, as described in A. Luzzago et al., Mimicking of discontinuous epitopes by phage displayed peptides, I. Epitope mapping of human H Ferritin using a phage library of constrained peptides, Gene (1993), 128, 51-57.

Sera from patients clinically characterized by being infected with the hepatitis C virus were then used.

The antibodies contained in these sera were immobilized on a solid matrix and incubated with the phage libraries. The phages specifically retained by these antibodies were then eluted and amplified.

Bacteria infected by the phages selected as above were plated on a solid culture medium and transferred onto nitrocellulose filters. Subsequently, these filters were incubated with serums from patients infected with HCV, different from the serum used for the initial enrichment. The phages specifically recognized by antibodies present in these sera were identified and isolated.

The phages identified as above were counter-selected for reactivity to antibodies present in the sera of individuals not infected with HCV, using the ELISA test.

The reaction specificity of the above phages with anti-HCV antibodies is evaluated statistically as follows. The frequency with which a significant number of sera from patients infected with HCV recognize the phage clones is determined in an ELISA test. In parallel, again using the ELISA test, absence of recognition of the same phage clones by a significant number of sera from individuals not infected with HCV is tested. For each phage clone, the specificity is evaluated by comparison of the frequency of recognition by sera from patients infected with HCV with that of

- 17 -

sera from non-infected individuals.

The epitope peptidic sequence identified in this manner was subsequently determined (SEQ ID from NO: 9 to NO: 30).

#### EXAMPLE 5

#### Demonstration of the effectiveness of the selected phages as diagnostic reagents for determination of the presence of anti-HCV antibodies in serum.

A process comprising the following operations was used:

40 human sera from individuals not infected with HCV were analyzed in an ELISA test for their ability to recognize specifically the phage clones, identified as described in example 4, displaying the peptide sequences described (SEQ ID from NO: 9 to NO: 23). The results show that none of the sera recognizes the sequences selected.

42 human sera from patients infected with HCV were analyzed in an ELISA test for their ability to recognize specifically the phage clones, identified as described in example 4, displaying the peptide sequences described (SEQ ID from NO: 9 to NO: 23). The results show that each phage clone is recognized specifically by the sera selected, with a frequency varying from 15 to 65% (see table I). The same results also show that each of the 42 sera tested specifically recognizes at least one of the sequences selected.

The sequences selected give an effective indication of the presence of anti-HCV antibodies in the blood of all patients examined, and do not react with blood from non-infected individuals. These data show that the group of phage clones selected forms a reliable system for the diagnosis of infection by the hepatitis C virus.

TABLE 1

Seq. Id	c1	c2	c3	c4	c5	c6	c7	c8	c9	c10	c11	c12	c13	c14	c15	c16	c17	c18	c19	c20	c21	Sequence
9																						LPAHGPSLS
10																						LPWGVARR
11																						PTHYTTSAP
12																						PTHYISSRH
13																						PTHYISTSL
14																						TRHYLRPGL
15																						PSHYVPRIY
16																						PPHLTSSCR
17																						KLNSRGSIS
18																						GKFPKSKPS
19																						FPGGPPLRA
20																						APSLPAGYL
21																						VPQSRLEPW
22																						NKREWAPPP
23																						NKTKQNPNL
Seq. Id	c22	c23	c24	c25	c26	c27	c28	c29	c30	c31	c32	c33	c34	c35	c36	c37	c38	c39	c40	c41	c42	Sequence
9																						LPAHGPSLS
10																						LPWGVARR
11																						PTHYTTSAP
12																						PTHYISSRH
13																						PTHYISTSL
14																						TRHYLRPGL
15																						PSHYVPRIY
16																						PPHLTSSCR
17																						KLNSRGSIS
18																						GKFPKSKPS
19																						FPGGPPLRA
20																						APSLPAGYL
21																						VPQSRLEPW
22																						NKREWAPPP
23																						NKTKQNPNL



" = Positive"



" = Negative"



" = Not analyzed"

- 19 -

Example 6Process for the preparation of specific diagnostic réagents and immunogens for the disease type II Cryoglobulinemia caused by and/or associated with the human Hepatitis C virus (HCV)

The following process was used according to the invention:

A first "library" of epitopes was prepared, made up of phages displaying oligopeptides with 9 amino acids on the surface of the capsid, expressed by random sequence oligonucleotidic inserts introduced into the gene coding for the protein VIII of the phage capsid. For expression of the recombinant proteins containing the epitope, the plasmid pC89 was used, engineered as described in F. Felici et al., Selection of Antibody Ligands from a Large Library of Oligopeptides Expressed on a Multivalent Exposition Vector, J. Mol. Biol. (1991), 222, 301-310, of which a portion of the genetic map is shown in the figure.

A second epitope "library" was prepared in the same way as the first, with the difference that two cysteine residues are present at the two ends of the insert, as described in A. Luzzago et al., Mimicking of discontinuous epitopes by phage displayed peptides, I. Epitope mapping of human H Ferritin using a phage library of constrained peptides, Gene (1993), 128, 51-57.

Samples of antibodies from individuals suffering from type II Cryoglobulinemia caused by and/or associated with the human hepatitis C virus (HCV) were immobilized on a solid matrix and incubated with the phage libraries. The phages specifically retained by said antibody were then eluted and amplified.

Bacteria infected by the phages selected as above were plated on a solid culture medium and transferred onto nitrocellulose filters. Subsequently, these filters were incubated with sera from patients

- 20 -

suffering from type II Cryoglobulinemia caused by and/or associated with human hepatitis C virus (HCV), different from the serum used for the first enrichment. The phages specifically recognized by antibodies present in these serums were identified and isolated.

The phages identified as above were counter-selected for reactivity to antibodies present in the serums of individuals not infected with type II Cryoglobulinemia caused by and/or associated with human hepatitis C virus (HCV), using the ELISA test.

The reaction specificity of the phages resulting from this counter-selection with antibodies specific for Type II Cryoglobulinemia caused by and/or associated with the human hepatitis C virus is evaluated statistically as follows. The frequency with which a significant number of sera from patients suffering from type II Cryoglobulinemia caused by and/or associated with the human hepatitis C virus (HCV) recognize the phage clones is determined in an ELISA test. In parallel, again using the ELISA test, absence of recognition of the same phage clones by a significant number of sera from individuals not suffering from the disease is tested. For each phage clone, the specificity is evaluated by comparison of the frequency of recognition by sera from patients suffering from the disease with that of sera from healthy individuals.

The epitope peptidic sequence identified in this manner was subsequently determined (SEQ ID from NO: 31 to NO: 42).

#### EXAMPLE 7

Demonstration of the effectiveness of the selected phages as diagnostic reagents for determination of the presence in the serum of antibodies specific for the disease type II Cryoglobulinemia caused by and/or associated with the human hepatitis C virus (HCV).

A process comprising the following operations was

- 21 -

used:

16 human sera from individuals not suffering from type II Cryoglobulinemia, caused by and/or associated with the human hepatitis C virus (HCV), were analyzed in an ELISA test for their ability to recognize specifically the phage clones, identified as described in example 6, showing the peptide sequences described (SEQ ID from NO: 31 to NO: 42). The results show that none of the serums recognizes the sequences selected.

80 human sera from patients suffering from type II Cryoglobulinemia, caused by and/or associated with the human hepatitis C virus (HCV), were analyzed in an ELISA test for their ability to recognize specifically the phage clones, identified as described in example 6, displaying the peptide sequences described (SEQ ID from NO: 31 to NO: 42). The results show that each phage clone is recognized specifically by the sera selected, with a frequency varying from 15 to 60%.

The sequences selected give an effective indication of the presence of antibodies specific for type II Cryoglobulinemia, caused by and/or associated with the human hepatitis C virus (HCV), in the blood of all patients examined, and do not react with blood from individuals not suffering from the disease. These data show that the group of phage clones selected forms a reliable system for diagnosis of the disease type II Cryoglobulinemia caused by and/or associated with the human hepatitis C virus (HCV).

The sequences selected show significant homology with the human LAG-3 gene sequence, which is specifically expressed by T lymphocytes and "natural killer" cells. The same antibodies used to select the phage clones react with the portion of the protein LAG-3 that is homologous with the sequence of the selected phage clones. This recognition can be abolished by competition with the selected phage clones. These results indicate that the selected phage clones can be

- 22 -

used to block the autoantibody binding to the protein LAG-3.

#### Example 8

Process for the preparation of diagnostic reagents for Type I Diabetes autoimmune disease and demonstration of the effectiveness of the selected phages as diagnostic reagents for determination of the presence in the blood of autoantibodies characterizing autoimmune diseases.

The following process was used according to the invention:

A first "library" of epitopes was prepared, made up of phages having oligopeptides with 9 amino acids displayed on the surface of the capsid, expressed by random sequence oligonucleotidic inserts introduced into the gene coding for the protein VIII of the phage capsid. For expression of the recombinant proteins containing the epitope, the plasmid pC89 was used, engineered as described in F. Felici et al., Selection of Antibody Ligands from a Large Library of Oligopeptides Expressed on a Multivalent Exposition Vector, J. Mol. Biol. (1991), 222, 301-310.

A second epitope "library" was prepared in the same way as the first, with the difference that two cysteine residues are present at the two ends of the insert, as described in A. Luzzago et al., Mimicking of discontinuous epitopes by phage displayed peptides, I. Epitope mapping of human H. Ferritin using a phage library of constrained peptides, Gene (1993), 128, 51-57.

A sample of sera from diabetic patients at the beginning of their type I diabetes clinical history was tested for the presence of antibodies specific for certain known markers characteristic of the disease, such as anti-insulin and ICA antibodies, or antibodies against the pancreatic islets. A serum with a high antibody content was then used for selection.

The antibodies contained in this serum were



- 23 -

immobilized on a solid matrix and incubated with the phage libraries. The phages specifically retained by these antibodies were then eluted and amplified.

Bacteria infected by the phages selected as above were plated on a solid culture medium and transferred onto nitrocellulose filters. Subsequently, these filters were incubated with sera from diabetic individuals different from the serum used for the first enrichment. The phages specifically recognized by antibodies present in these sera were identified and isolated.

The phages identified as above were counter-selected for reactivity to antibodies present in the sera of individuals not suffering from type I diabetes, using the ELISA test.

The phages resulting from this counter-selection were checked for specificity of reaction by means of large scale screening using sera from diabetic patients at different stages of the disease. The nucleotidic sequence coding for the epitopes identified in this manner was then determined.

The phage clones were analyzed using the following process:

25 human sera taken from individuals at the time of the first clinical appearance of type I diabetes were analyzed using an ELISA test for their ability to recognize specifically the phage clones, identified as described and displaying the peptide sequences described (SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47). The results show that 36% of the sera recognized specifically at least one of the 5 sequences selected.

16 human sera taken from individuals with autoimmune pathologies other than diabetes were analyzed using an ELISA test for their ability to recognize specifically the phage clones, identified as described in example 8, and displaying the peptide

- 24 -

sequences described (SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47). The results show that 18% of the sera recognized specifically at least one of the 5 sequences selected.

50 human sera taken from individuals not suffering from diabetes or from other autoimmune pathologies were analyzed using an ELISA test for their ability to recognize specifically the phage clones, identified as described in example 8, and displaying the peptide sequences described (SEQ ID NO: 43, SEQ ID NO: 44, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 47). The results show that none of the sera recognized specifically any of the 5 sequences selected.

#### Example 9

Process for the preparation of diagnostic reagents and immunogens specific for detection and induction of the immune response of the human tumoral protein NEU (p185<sup>HER2</sup>)

The following process was used according to the invention:

The monoclonal antibodies MGr2 and MGr6 were identified as recognizing specifically the human tumoral protein NEU (p185<sup>HER2</sup>).

Two epitope "libraries" were prepared, made up of phages displaying oligopeptides on the surface of the capsid (as described in example 1), which were then examined using the biopanning method (V. Parmley, S.F. & Smith, G.P., Antibody-selectable filamentous fd phage vectors: affinity purification of target genes, Gene (1988) 73, 305-318) to select the clones that react specifically with the above mentioned antibodies.

Bacteria infected by the phages selected as above were plated on a solid culture medium and transferred onto nitrocellulose filters. Subsequently, these filters were incubated with the monoclonal antibodies MGr2 and MGr6. The reaction specificity of the phages identified in this way was controlled using the ELISA

- 25 -

test.

The peptidic sequence of the epitopes identified as above was subsequently determined by means of analysis of the corresponding nucleotidic coding sequence (for MGr2 SEQ ID from NO: 48 to NO: 56, for MGr6 SEQ ID from NO: 57 to NO: 68).

The phages, selected as above, were amplified, purified and injected into test animals according to the procedure described in Example 2. The serum from animals immunized in this manner was tested for the presence of antibodies capable of interacting with NEU, using an immunohistochemical test.

This method underlined the presence of a significant level of anti-NEU antibodies in the blood of test animals immunized in this manner.

The results obtained with the phages described in this example demonstrate the effectiveness of said phages as antigenic and immunogenic substituted of the human tumoral protein NEU. This may allow the phages thus obtained, or derivatives thereof, to be used as reagents for the detection of anti-NEU antibodies in sera from patients suffering from tumors and/or as specific immunogens to stimulate an anti-NEU antibody response.

- 26 -

## SEQUENCE LISTING

## GENERAL INFORMATION

- (i) APPLICANT: ISTITUTO DI RICERCHE DI BIOLOGIA  
MOLECOLARE P. ANGELETTI S.p.A.
- (ii) TITLE OF INVENTION: Process for the preparation of immunogens and diagnostic reagents, and immunogens and diagnostic reagents thereby obtained
- (iii) NUMBER OF SEQUENCES: 68
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Società Italiana Brevetti
  - (B) STREET: Piazza di Pietra, 39
  - (C) CITY: Rome
  - (D) COUNTRY: Italy
  - (E) POSTAL CODE: I-00186
- (viii) ATTORNEY INFORMATION
  - (A) NAME: DI CERBO, Mario (Dr.)
  - (B) REFERENCE: RM/090041/MDC
- (ix) TELECOMMUNICATION INFORMATION
  - (A) TELEPHONE: 06/6785941
  - (B) TELEFAX: 06/6794692
  - (C) TELEX: 612287 ROPAT
- (1) INFORMATION FOR SEQ ID NO: 1
  - (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 15 amino-acids
    - (B) TYPE: aminoacidic
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: recombinant protein
  - (iii) HYPOTHETICAL: yes
  - (v) FRAGMENT TYPE: internal
  - (vii) IMMEDIATE SOURCE
    - (A) LIBRARY: of recombinant peptides on phage
    - (B) CLONE: phagic
  - (ix) FURTHER CHARACTERISTICS
    - (A) NAME: polypeptide
    - (B) IDENTIFICATION METHOD: selection with

- 27 -

## specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1

Glu	Phe	Cys	Arg	Thr	Cys	Ala	His	Pro	Gly	Glu	His	Ala	Gly	Asp
1				5					10					15

(2) INFORMATION FOR SEQ ID NO: 2

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2

Glu	Phe	Cys	Gly	Pro	Phe	Tyr	Leu	Ser	Ala	Pro	Gln	Cys	Gly	Asp
1				5					10					15

(3) INFORMATION FOR SEQ ID NO: 3

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

- 28 -

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3

Glu Phe Cys Gly Pro Phe Phe Leu Ala Ala Ser Val Cys Gly Asp  
1                      5                      10                      15

(4) INFORMATION FOR SEQ ID NO: 4

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 47 pairs of bases

(B) TYPE: nucleotidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(ix) FURTHER CHARACTERISTICS

(A) NAME: polynucleotide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4

GAATTCTGCC GAACCTGCGC CCATCCAGGT GAGCATGCCG GGGATCC 47

(5) INFORMATION FOR SEQ ID NO: 5

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 47 pairs of bases

(B) TYPE: nucleotidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(ix) FURTHER CHARACTERISTICS

(A) NAME: polynucleotide

- 29 -

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5

GAATTCTGCG GGCCTTTTTA TCTCTCTGCA CCTCAGTGCG GGGATCC 47

(6) INFORMATION FOR SEQ ID NO: 6

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 47 pairs of bases

(B) TYPE: nucleotidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(ix) FURTHER CHARACTERISTICS

(A) NAME: polynucleotide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6

GAATTCTGCG GTCCCTTCTT TCTCGCGGCT TCCGTATGCG GGGATCC 47

#### SEQUENCE LISTING

##### GENERAL INFORMATION

(i) APPLICANT: ISTITUTO DI RICERCHE DI BIOLOGIA  
MOLECOLARE P. ANGELETTI S.p.A.

(ii) TITLE OF INVENTION: Process for the prepara-  
tion of immunogens and diagnostic reagents,  
and immunogens and diagnostic reagents thereby  
obtained

(iii) NUMBER OF SEQUENCES: 68

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Società Italiana Brevetti

(B) STREET: Piazza di Pietra, 39

(C) CITY: Rome

(D) COUNTRY: Italy

(E) POSTAL CODE: I-00186

- 30 -

## (viii) ATTORNEY INFORMATION

(A) NAME: DI CERBO, Mario (Dr.)

(B) REFERENCE: RM/090041/MDC

## (ix) TELECOMMUNICATION INFORMATION

(A) TELEPHONE: 06/6785941

(B) TELEFAX: 06/6794692

(C) TELEX: 612287 ROPAT

## (1) INFORMATION FOR SEQ ID NO: 1

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

## (vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1

Glu Phe Cys Arg Thr Cys Ala His Pro Gly Glu His Ala Gly Asp  
1                      5                      10                      15

## (2) INFORMATION FOR SEQ ID NO: 2

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

## (vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage



- 31 -

(B) CLONE: phagic

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2

Glu	Phe	Cys	Gly	Pro	Phe	Tyr	Leu	Ser	Ala	Pro	Gln	Cys	Gly	Asp
1				5					10					15

## (3) INFORMATION FOR SEQ ID NO: 3

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

## (vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3

Glu	Phe	Cys	Gly	Pro	Phe	Phe	Leu	Ala	Ala	Ser	Val	Cys	Gly	Asp
1				5					10					15

## (4) INFORMATION FOR SEQ ID NO: 4

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 47 pairs of bases

(B) TYPE: nucleotidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

(v) FRAGMENT TYPE: internal

- 32 -

- (vii) IMMEDIATE SOURCE
    - (A) LIBRARY: of recombinant peptides on phage
  - (ix) FURTHER CHARACTERISTICS
    - (A) NAME: polynucleotide
    - (B) IDENTIFICATION METHOD: selection with specific antibodies
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4  
GAATTCTGCC GAACCTGCGC CCATCCAGGT GAGCATGCGG GGGATCC 47
- (5) INFORMATION FOR SEQ ID NO: 5
- (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 47 pairs of bases
    - (B) TYPE: nucleotidic
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA
  - (iii) HYPOTHETICAL: no
  - (iv) ANTISENSE: no
  - (v) FRAGMENT TYPE: internal
  - (vii) IMMEDIATE SOURCE
    - (A) LIBRARY: of recombinant peptides on phage
  - (ix) FURTHER CHARACTERISTICS
    - (A) NAME: polynucleotide
    - (B) IDENTIFICATION METHOD: selection with specific antibodies
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5  
GAATTCTGCG GGCCTTTTTA TCTCTGCA CCTCAGTGCG GGGATCC 47
- (6) INFORMATION FOR SEQ ID NO: 6
- (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 47 pairs of bases
    - (B) TYPE: nucleotidic
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA
  - (iii) HYPOTHETICAL: no
  - (iv) ANTISENSE: no
  - (v) FRAGMENT TYPE: internal
  - (vii) IMMEDIATE SOURCE

- 33 -

- (A) LIBRARY: of recombinant peptides on phage
- (ix) FURTHER CHARACTERISTICS
- (A) NAME: polynucleotide
- (B) IDENTIFICATION METHOD: selection with  
specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6
- GAATTCTGCG GTCCCTTCTT TCTCGCGGCT TCCGTATGCG GGGATCC 47

## SEQUENCE LISTING

## GENERAL INFORMATION

- (i) APPLICANT: ISTITUTO DI RICERCHE DI BIOLOGIA  
MOLECOLARE P. ANGELETTI S.p.A.
- (ii) TITLE OF INVENTION: Process for the prepara-  
tion of immunogens and diagnostic reagents,  
and immunogens and diagnostic reagents thereby  
obtained
- (iii) NUMBER OF SEQUENCES: 68
- (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: Società Italiana Brevetti
- (B) STREET: Piazza di Pietra, 39
- (C) CITY: Rome
- (D) COUNTRY: Italy
- (E) POSTAL CODE: I-00186
- (viii) ATTORNEY INFORMATION
- (A) NAME: DI CERBO, Mario (Dr.)
- (B) REFERENCE: RM/090041/MDC
- (ix) TELECOMMUNICATION INFORMATION
- (A) TELEPHONE: 06/6785941
- (B) TELEFAX: 06/6794692
- (C) TELEX: 612287 ROPAT
- (1) INFORMATION FOR SEQ ID NO: 1
- (i) SEQUENCE CHARACTERISTICS
- (A) LENGTH: 15 amino-acids
- (B) TYPE: aminoacidic
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes

- 34 -

- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
- (A) LIBRARY: of recombinant peptides on phage
- (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
- (A) NAME: polypeptide
- (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1
- |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Phe | Cys | Arg | Thr | Cys | Ala | His | Pro | Gly | Glu | His | Ala | Gly | Asp |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |
- (2) INFORMATION FOR SEQ ID NO: 2
- (i) SEQUENCE CHARACTERISTICS
- (A) LENGTH: 15 amino-acids
- (B) TYPE: aminoacidic
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
- (A) LIBRARY: of recombinant peptides on phage
- (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
- (A) NAME: polypeptide
- (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2
- |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Phe | Cys | Gly | Pro | Phe | Tyr | Leu | Ser | Ala | Pro | Gln | Cys | Gly | Asp |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |
- (3) INFORMATION FOR SEQ ID NO: 3
- (i) SEQUENCE CHARACTERISTICS
- (A) LENGTH: 15 amino-acids
- (B) TYPE: aminoacidic
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein

- 35 -

- (iii) HYPOTHETICAL: yes  
(v) FRAGMENT TYPE: internal  
(vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic  
(ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with specific antibodies  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3  
Glu Phe Cys Gly Pro Phe Phe Leu Ala Ala Ser Val Cys Gly Asp  
1 5 10 15  
(4) INFORMATION FOR SEQ ID NO: 4  
(i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 47 pairs of bases  
(B) TYPE: nucleotidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(ii) MOLECULE TYPE: DNA  
(iii) HYPOTHETICAL: no  
(iv) ANTISENSE: no  
(v) FRAGMENT TYPE: internal  
(vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(ix) FURTHER CHARACTERISTICS  
(A) NAME: polynucleotide  
(B) IDENTIFICATION METHOD: selection with specific antibodies  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4  
GAATTCTGCC GAACCTGCGC CCATCCAGGT GAGCATGCGG GGGATCC 47  
(5) INFORMATION FOR SEQ ID NO: 5  
(i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 47 pairs of bases  
(B) TYPE: nucleotidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(ii) MOLECULE TYPE: DNA

- 36 -

- (iii) HYPOTHETICAL: no
- (iv) ANTISENSE: no
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polynucleotide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5  
GAATTCTGCG GGCCTTTTTA TCTCTCTGCA CCTCAGTGCG GGGATCC 47
- (6) INFORMATION FOR SEQ ID NO: 6
  - (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 47 pairs of bases
    - (B) TYPE: nucleotidic
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA
  - (iii) HYPOTHETICAL: no
  - (iv) ANTISENSE: no
  - (v) FRAGMENT TYPE: internal
  - (vii) IMMEDIATE SOURCE
    - (A) LIBRARY: of recombinant peptides on phage
  - (ix) FURTHER CHARACTERISTICS
    - (A) NAME: polynucleotide
    - (B) IDENTIFICATION METHOD: selection with specific antibodies
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6  
GAATTCTGCG GTCCCTTCTT TCTCGCGGCT TCCGTATGCG GGGATCC 47

## SEQUENCE LISTING

## GENERAL INFORMATION

- (i) APPLICANT: ISTITUTO DI RICERCHE DI BIOLOGIA  
MOLECOLARE P. ANGELETTI S.p.A.
- (ii) TITLE OF INVENTION: Process for the preparation of immunogens and diagnostic reagents, and immunogens and diagnostic reagents thereby obtained

- 37 -

- (iii) NUMBER OF SEQUENCES: 68
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Società Italiana Brevetti
  - (B) STREET: Piazza di Pietra, 39
  - (C) CITY: Rome
  - (D) COUNTRY: Italy
  - (E) POSTAL CODE: I-00186
- (viii) ATTORNEY INFORMATION
  - (A) NAME: DI CERBO, Mario (Dr.)
  - (B) REFERENCE: RM/090041/MDC
- (ix) TELECOMMUNICATION INFORMATION
  - (A) TELEPHONE: 06/6785941
  - (B) TELEFAX: 06/6794692
  - (C) TELEX: 612287 ROPAT
- (1) INFORMATION FOR SEQ ID NO: 1
  - (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 15 amino-acids
    - (B) TYPE: aminoacidic
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: recombinant protein
  - (iii) HYPOTHETICAL: yes
  - (v) FRAGMENT TYPE: internal
  - (vii) IMMEDIATE SOURCE
    - (A) LIBRARY: of recombinant peptides on phage
    - (B) CLONE: phagic
  - (ix) FURTHER CHARACTERISTICS
    - (A) NAME: polypeptide
    - (B) IDENTIFICATION METHOD: selection with specific antibodies
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1

Glu	Phe	Cys	Arg	Thr	Cys	Ala	His	Pro	Gly	Glu	His	Ala	Gly	Asp
1				5					10					15
- (2) INFORMATION FOR SEQ ID NO: 2
  - (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 15 amino-acids
    - (B) TYPE: aminoacidic

- 38 -

- (C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
- (A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
- (A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2
- Glu Phe Cys Gly Pro Phe Tyr Leu Ser Ala Pro Gln Cys Gly Asp  
1 5 10 15
- (3) INFORMATION FOR SEQ ID NO: 3
- (i) SEQUENCE CHARACTERISTICS
- (A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
- (A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
- (A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3
- Glu Phe Cys Gly Pro Phe Phe Leu Ala Ala Ser Val Cys Gly Asp  
1 5 10 15
- (4) INFORMATION FOR SEQ ID NO: 4
- (i) SEQUENCE CHARACTERISTICS
- (A) LENGTH: 47 pairs of bases



- 39 -

- (B) TYPE: nucleotidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA
  - (iii) HYPOTHETICAL: no
  - (iv) ANTISENSE: no
  - (v) FRAGMENT TYPE: internal
  - (vii) IMMEDIATE SOURCE
    - (A) LIBRARY: of recombinant peptides on phage
  - (ix) FURTHER CHARACTERISTICS
    - (A) NAME: polynucleotide
    - (B) IDENTIFICATION METHOD: selection with specific antibodies
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4  
GAATTCTGCC GAACCTGCGC CCATCCAGGT GAGCATGCGG GGGATCC 47
- (5) INFORMATION FOR SEQ ID NO: 5
- (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 47 pairs of bases
    - (B) TYPE: nucleotidic
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA
  - (iii) HYPOTHETICAL: no
  - (iv) ANTISENSE: no
  - (v) FRAGMENT TYPE: internal
  - (vii) IMMEDIATE SOURCE
    - (A) LIBRARY: of recombinant peptides on phage
  - (ix) FURTHER CHARACTERISTICS
    - (A) NAME: polynucleotide
    - (B) IDENTIFICATION METHOD: selection with specific antibodies
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5  
GAATTCTGCG GGCCTTTTTA TCTCTCTGCA CCTCAGTGCG GGGATCC 47
- (6) INFORMATION FOR SEQ ID NO: 6
- (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 47 pairs of bases
    - (B) TYPE: nucleotidic

- 40 -

- (C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(ii) MOLECULE TYPE: DNA  
(iii) HYPOTHETICAL: no  
(iv) ANTISENSE: no  
(v) FRAGMENT TYPE: internal  
(vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(ix) FURTHER CHARACTERISTICS  
(A) NAME: polynucleotide  
(B) IDENTIFICATION METHOD: selection with specific antibodies  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6  
GAATTCTGCG GTCCCTTCTT TCTCGGGCT TCCGTATGCG GGGATCC 47  
(7) INFORMATION FOR SEQ ID NO: 7  
(i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(ii) MOLECULE TYPE: recombinant protein  
(iii) HYPOTHETICAL: yes  
(v) FRAGMENT TYPE: internal  
(vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic  
(ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with specific antibodies  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7  
Glu Phe Cys Gly Pro Phe Phe Leu Ser Pro Thr Ser Cys Gly Asp  
1 5 10 15  
(8) INFORMATION FOR SEQ ID NO: 8  
(i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 47 pairs of bases  
(B) TYPE: nucleotidic

- 41 -

- (C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (iii) HYPOTHETICAL: no
- (iv) ANTISENSE: no
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage
- (ix) FURTHER CHARACTERISTICS  
(A) NAME: polynucleotide  
(B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8  
GAATTCTGCG GTCCGTTTTT TCTCTCCCG ACGTCATGCG GGGATCC 47
- (9) INFORMATION FOR SEQ ID NO: 9
- (i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9  
Glu Phe Cys Leu Pro Ala His Gly Pro Ser Leu Ser Cys Gly Asp  
1 5 10 15
- (10) INFORMATION FOR SEQ ID NO: 10
- (i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic

- 42 -

- (C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10
- |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Phe | Cys | Leu | Pro | Trp | Gly | Val | Ala | Ala | Arg | Arg | Cys | Gly | Asp |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |
- (11) INFORMATION FOR SEQ ID NO: 11
- (i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11
- |     |     |     |     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Glu | Phe | Cys | Pro | Thr | His | Tyr | Thr | Thr | Ser | Ala | Pro | Cys | Gly | Asp |
| 1   |     |     |     | 5   |     |     |     |     | 10  |     |     |     |     | 15  |
- (12) INFORMATION FOR SEQ ID NO: 12
- (i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids

- 43 -

- (B) TYPE: aminoacidic
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12  
Glu Phe Cys Pro Thr His Tyr Ile Ser Ser Arg His Cys Gly Asp  
1                      5                      10                      15
- (13) INFORMATION FOR SEQ ID NO: 13
  - (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 15 amino-acids
    - (B) TYPE: aminoacidic
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: recombinant protein
  - (iii) HYPOTHETICAL: yes
  - (v) FRAGMENT TYPE: internal
  - (vii) IMMEDIATE SOURCE
    - (A) LIBRARY: of recombinant peptides on phage
    - (B) CLONE: phagic
  - (ix) FURTHER CHARACTERISTICS
    - (A) NAME: polypeptide
    - (B) IDENTIFICATION METHOD: selection with specific antibodies
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13  
Glu Phe Cys Pro Thr His Tyr Ile Ser Thr Ser Leu Cys Gly Asp  
1                      5                      10                      15
- (14) INFORMATION FOR SEQ ID NO: 14
  - (i) SEQUENCE CHARACTERISTICS

- 44 -

- (A) LENGTH: 15 amino-acids
- (B) TYPE: aminoacidic
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14  
Glu Phe Cys Thr Arg His Tyr Leu Arg Pro Gly Leu Cys Gly Asp  
1                      5                      10                      15
- (15) INFORMATION FOR SEQ ID NO: 15
  - (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 15 amino-acids
    - (B) TYPE: aminoacidic
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: recombinant protein
  - (iii) HYPOTHETICAL: yes
  - (v) FRAGMENT TYPE: internal
  - (vii) IMMEDIATE SOURCE
    - (A) LIBRARY: of recombinant peptides on phage
    - (B) CLONE: phagic
  - (ix) FURTHER CHARACTERISTICS
    - (A) NAME: polypeptide
    - (B) IDENTIFICATION METHOD: selection with specific antibodies
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15  
Glu Phe Cys Pro Ser His Tyr Val Pro Arg Ile Tyr Cys Gly Asp  
1                      5                      10                      15
- (16) INFORMATION FOR SEQ ID NO: 16

- 45 -

- (i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with  
specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16
- Glu Phe Cys Pro Pro His Leu Thr Leu Ser Ser Cys Arg Gly Asp  
1 5 10 15
- (17) INFORMATION FOR SEQ ID NO: 17
- (i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with  
specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17
- Glu Phe Cys Lys Leu Asn Ser Arg Gly Ser Ile Ser Cys Gly Asp  
1 5 10 15

- 46 -

## (18) INFORMATION FOR SEQ ID NO: 18

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18

Glu Phe Cys Gly Lys Phe Pro Gly Ser Lys Pro Ser Cys Gly Asp  
1                      5                      10                      15

## (19) INFORMATION FOR SEQ ID NO: 19

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19

Glu Phe Cys Phe Pro Gly Gly Pro Pro Leu Arg Ala Cys Gly Asp



**1                      5                      10                      15**

### (i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 15 amino-acids
- (B) TYPE: aminoacidic
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20

Glu Phe Cys Ala Pro Ser Leu Pro Ala Gly Tyr Leu Cys Gly Asp

**1                      5                      10                      15**

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 15 amino-acids
- (B) TYPE: aminoacidic
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21

- 48 -

Glu Phe Cys Gln Val Pro Gln Ser Arg Leu Glu Pro Trp Gly Asp  
1 5 10 15

(22) INFORMATION FOR SEQ ID NO: 22

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 12 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22

Glu Phe Asn Lys Arg Glu Trp Ala Pro Pro Pro Asp  
1                5                      10          12

(23) INFORMATION FOR SEQ ID NO: 23

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 12 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies

- 49 -

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23

Glu Phe Asn Lys Thr Lys Gln Asn Pro Asn Leu Asp  
1                      5                      10              12

## (24) INFORMATION FOR SEQ ID NO: 24

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 12 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24

Glu Phe Thr Ser Leu Gln Pro Asp Arg Ala Gln Asp  
1                      5                      10              12

## (25) INFORMATION FOR SEQ ID NO: 25

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 12 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with

- 50 -

specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25

Glu Phe Ser Gly Leu Arg Pro Gly Lys Phe Gln Asp  
1                      5                      10              12

## (26) INFORMATION FOR SEQ ID NO: 26

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 12 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26

Glu Phe Thr Gly Val Arg Glu Ile Ser Phe Gly Asp  
1                      5                      10              12

## (27) INFORMATION FOR SEQ ID NO: 27

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 12 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

- 51 -

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27

Glu Phe Thr Gly Leu Arg Glu Ser Pro Ser Met Asp  
1                      5                      10              12

(28) INFORMATION FOR SEQ ID NO: 28

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 12 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28

Glu Phe Ser His Pro His Phe Ser Gly Leu Glu Asp  
1                      5                      10              12

(29) INFORMATION FOR SEQ ID NO: 29

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 12 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

- 52 -

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29

Glu Phe Thr Gly Leu Arg Ser Arg Tyr Pro Ala Asp

1                      5                      10              12

(30) INFORMATION FOR SEQ ID NO: 30

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 12 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30

Glu Phe Thr Gly Leu Arg His Lys Thr Ser Ala Asp

1                      5                      10              12

(31) INFORMATION FOR SEQ ID NO: 31

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

- 53 -

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31

Glu	Phe	Cys	His	Pro	Leu	Ala	Pro	Ala	Gly	Thr	Phe	Cys	Gly	Asp
1				5					10					15

## (32) INFORMATION FOR SEQ ID NO: 32

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 14 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32

Glu	Phe	Cys	His	Pro	Leu	Ala	Pro	Leu	Asn	Phe	Cys	Gly	Asp
1				5				10					14

## (33) INFORMATION FOR SEQ ID NO: 33

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

- 54 -

(B) CLONE: phagic

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33

Glu	Phe	Cys	Gly	His	Pro	Leu	Ala	Pro	Pro	Gln	Ala	Cys	Gly	Asp
1				5				10					15	

## (34) INFORMATION FOR SEQ ID NO: 34

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

## (vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34

Glu	Phe	Cys	His	Pro	Leu	Ser	Pro	Arg	Pro	Leu	Gln	Cys	Gly	Asp
1				5				10					15	

## (35) INFORMATION FOR SEQ ID NO: 35

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

## (vii) IMMEDIATE SOURCE



- 55 -

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35

Glu	Phe	Cys	His	Pro	Leu	Ala	Pro	Pro	His	Pro	Ser	Cys	Gly	Asp
1				5				10					15	

## (36) INFORMATION FOR SEQ ID NO: 36

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

## (vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36

Glu	Phe	Cys	His	Pro	Leu	Ser	Pro	His	Pro	Ser	Tyr	Cys	Gly	Asp
1				5				10					15	

## (37) INFORMATION FOR SEQ ID NO: 37

## (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

- 56 -

- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic

- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37

Glu Phe Cys His Pro Leu Ala Thr Gly Pro His Leu Cys Gly Asp  
1                      5                                      10                                      15

- (38) INFORMATION FOR SEQ ID NO: 38

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE

- (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic

- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38

Glu Phe Cys His Pro Leu Ala Pro Ser Pro Ala Val Cys Gly Asp  
1                      5                                      10                                      15

- (39) INFORMATION FOR SEQ ID NO: 39

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes

- 57 -

(v) FRAGMENT TYPE: internal  
(vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic  
(ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with  
specific antibodies  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39  
Glu Phe Cys His Pro Leu Pro Pro Ala Ala Thr Phe Cys Gly Asp  
1 5 10 15

(40) INFORMATION FOR SEQ ID NO: 40  
(i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(ii) MOLECULE TYPE: recombinant protein  
(iii) HYPOTHETICAL: yes  
(v) FRAGMENT TYPE: internal  
(vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic  
(ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with  
specific antibodies  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40

Glu Phe Cys His Pro Leu Ala Pro Val Pro Arg Gln Cys Gly Asp  
1 5 10 15

(41) INFORMATION FOR SEQ ID NO: 41  
(i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41

(42) INFORMATION FOR SEQ ID NO: 42

Glu Phe Cys Asn His Pro Leu Ser Pro Ser Gly Ala Cys Gly Asp  
1                      5                      10                      15

(43) INFORMATION FOR SEQ ID NO: 43

(i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic  
(C) STRANDEDNESS: single

- 59 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43

Glu	Phe	Cys	His	Ala	Val	Lys	Gly	Phe	Ser	Ala	Val	Cys	Gly	Asp
1				5				10					15	

## (44) INFORMATION FOR SEQ ID NO: 44

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44

Glu	Phe	Cys	His	Ala	Val	Lys	Val	Gly	Asn	Pro	Ser	Cys	Gly	Asp
1				5				10					15	

## (45) INFORMATION FOR SEQ ID NO: 45

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

- 60 -

(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(ii) MOLECULE TYPE: recombinant protein  
(iii) HYPOTHETICAL: yes  
(v) FRAGMENT TYPE: internal  
(vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic  
(ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with  
specific antibodies  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45  
Glu Phe Cys His Ala Thr Lys Thr Pro Trp Thr Thr Cys Gly Asp  
1 5 10 15

(46) INFORMATION FOR SEQ ID NO: 46  
(i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(ii) MOLECULE TYPE: recombinant protein  
(iii) HYPOTHETICAL: yes  
(v) FRAGMENT TYPE: internal  
(vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic  
(ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with  
specific antibodies  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46  
Glu Phe Cys Arg Ala Pro Ser Gly Val Ile Val Gln Cys Gly Asp  
1 5 10 15

(47) INFORMATION FOR SEQ ID NO: 47  
(i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids

- 61 -

- (B) TYPE: aminoacidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein  
(iii) HYPOTHETICAL: yes  
(v) FRAGMENT TYPE: internal  
(vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with  
specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47  
Glu Phe Cys Gly Gly Ala Ser Ser Gly Cys Lys Pro Cys Gly Asp  
1 5 10 15
- (48) INFORMATION FOR SEQ ID NO: 48
- (i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 15 amino-acids  
(B) TYPE: aminoacidic  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein  
(iii) HYPOTHETICAL: yes  
(v) FRAGMENT TYPE: internal  
(vii) IMMEDIATE SOURCE  
(A) LIBRARY: of recombinant peptides on phage  
(B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS  
(A) NAME: polypeptide  
(B) IDENTIFICATION METHOD: selection with  
specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48  
Gly Glu Phe Tyr Thr Pro Thr Trp Met Leu Pro Glu Asp Pro Lys  
1 5 10 15
- (49) INFORMATION FOR SEQ ID NO: 49
- (i) SEQUENCE CHARACTERISTICS

- 62 -

- (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: recombinant protein
  - (iii) HYPOTHETICAL: yes
  - (v) FRAGMENT TYPE: internal
  - (vii) IMMEDIATE SOURCE
    - (A) LIBRARY: of recombinant peptides on phage
    - (B) CLONE: phagic
  - (ix) FURTHER CHARACTERISTICS
    - (A) NAME: polypeptide
    - (B) IDENTIFICATION METHOD: selection with specific antibodies
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49
- Gly Glu Phe Ser Pro Pro Trp Met Leu Pro Ser Val Asp Pro Lys  
1                      5                      10                      15
- (50) INFORMATION FOR SEQ ID NO: 50
- (i) SEQUENCE CHARACTERISTICS
    - (A) LENGTH: 15 amino-acids
    - (B) TYPE: aminoacidic
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: recombinant protein
  - (iii) HYPOTHETICAL: yes
  - (v) FRAGMENT TYPE: internal
  - (vii) IMMEDIATE SOURCE
    - (A) LIBRARY: of recombinant peptides on phage
    - (B) CLONE: phagic
  - (ix) FURTHER CHARACTERISTICS
    - (A) NAME: polypeptide
    - (B) IDENTIFICATION METHOD: selection with specific antibodies
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50
- Gly Glu Phe Thr Pro Asn Trp Met Leu Gln Asn Leu Asp Pro Lys  
1                      5                      10                      15
- (51) INFORMATION FOR SEQ ID NO: 51



- 63 -

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51

Gly Glu Phe Thr Pro Arg Trp Met Leu Ser Arg Glu Asp Pro Lys  
1                      5                      10                      15

(52) INFORMATION FOR SEQ ID NO: 52

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52

Gly Glu Phe Thr Pro Thr Trp Met Leu Ala Arg Trp Asp Pro Lys  
1                      5                      10                      15

- 64 -

## (53) INFORMATION FOR SEQ ID NO: 53

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53

Glu Phe Cys Gly Pro Leu Asp Ser Leu Phe Ala Gln Cys Gly Asp  
1                      5                      10                      15

## (54) INFORMATION FOR SEQ ID NO: 54

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54

Glu Phe Cys Gly Pro Ile Ser Ala Leu Phe Ala Ser Cys Gly Asp

- 65 -

**1                      5                      10                      15**

(55) INFORMATION FOR SEQ ID NO: 55

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55

Glu Phe Cys Gly Pro Ile His Ala Leu Phe Leu Asp Cys Gly Asp

**1                      5                      10                      15**

## (56) INFORMATION FOR SEQ ID NO: 56

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56

- 66 -

Glu Phe Cys Gly Pro Ile Ser Ser Leu Phe Gly Asp Cys Gly Asp  
1                      5                      10                      15

## (57) INFORMATION FOR SEQ ID NO: 57

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57

Gly Glu Phe Ile Cys His Ser Asp Cys Ala Ala Gly Asp Pro Lys  
1                      5                      10                      15

## (58) INFORMATION FOR SEQ ID NO: 58

- (i) SEQUENCE CHARACTERISTICS
  - (A) LENGTH: 15 amino-acids
  - (B) TYPE: aminoacidic
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: recombinant protein
- (iii) HYPOTHETICAL: yes
- (v) FRAGMENT TYPE: internal
- (vii) IMMEDIATE SOURCE
  - (A) LIBRARY: of recombinant peptides on phage
  - (B) CLONE: phagic
- (ix) FURTHER CHARACTERISTICS
  - (A) NAME: polypeptide
  - (B) IDENTIFICATION METHOD: selection with specific antibodies

- 67 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58

Gly Glu Phe Ile Cys His Ser Asp Cys Ala Ser Gly Asp Pro Lys  
1                      5                      10                      15

(59) INFORMATION FOR SEQ ID NO: 59

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59

Gly Glu Phe Ile Ala Cys His Ser Asp Cys Gly Ser Asp Pro Lys  
1                      5                      10                      15

(60) INFORMATION FOR SEQ ID NO: 60

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with

- 68 -

## specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60

Gly Glu Phe Trp Thr Pro Leu Lys Cys Asp Ala Leu Asp Pro Lys  
1                      5                      10                      15

(61) INFORMATION FOR SEQ ID NO: 61

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61

Gly Glu Phe Pro Ala Ala Phe Gly Lys Leu Gly Val Asp Pro Lys  
1                      5                      10                      15

(62) INFORMATION FOR SEQ ID NO: 62

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

- 69 -

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62

Glu	Phe	Cys	Leu	Val	Leu	Pro	Lys	Val	Lys	Met	Ala	Cys	Gly	Asp
1				5				10					15	

(63) INFORMATION FOR SEQ ID NO: 63

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63

Glu	Phe	Cys	Ala	Arg	Leu	Pro	Val	Leu	Lys	Leu	Val	Cys	Gly	Asp
1					5				10				15	

(64) INFORMATION FOR SEQ ID NO: 64

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

- 70 -

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64

Glu	Phe	Cys	Ile	Trp	Leu	Pro	Arg	Ile	Lys	Leu	Ser	Cys	Gly	Asp
1				5					10					15

(65) INFORMATION FOR SEQ ID NO: 65

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65

Glu	Phe	Cys	Phe	Ser	Ile	Pro	Ser	Leu	Lys	Thr	Val	Cys	Gly	Asp
1				5					10					15

(66) INFORMATION FOR SEQ ID NO: 66

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic



- 71 -

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66

Glu Phe Cys Phe Arg Gly Pro Arg Gln Lys Leu Tyr Cys Gly Asp  
1 5 10 15

(67) INFORMATION FOR SEQ ID NO: 67

### (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) **FRAGMENT TYPE:** internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

(B) CLONE: phagic

(ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with specific antibodies

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67

Glu Phe Cys Leu Pro Leu Leu Gly Arg Lys Thr Met Cys Gly Asp  
1                5                      10                          15

(68) INFORMATION FOR SEQ ID NO: 68

### (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15 amino-acids

(B) TYPE: aminoacidic

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: recombinant protein

(iii) HYPOTHETICAL: yes

(v) FRAGMENT TYPE: internal

(vii) IMMEDIATE SOURCE

(A) LIBRARY: of recombinant peptides on phage

- 72 -

(B) CLONE: phagic

## (ix) FURTHER CHARACTERISTICS

(A) NAME: polypeptide

(B) IDENTIFICATION METHOD: selection with  
specific antibodies

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 68

Glu	Phe	Cys	Trp	Ile	His	His	Leu	Leu	Lys	Val	Val	Cys	Gly	Asp
1					5				10				15	

- 73 -

CLAIMS

1. Process for the preparation of immunogens or diagnostic reagents that mimic an antigen or a pathogenic organism specific to a disease, essentially characterized by the following operations:

- identification of at least one antibody that reacts with the antigen or pathogenic organism specific to the disease;

- construction of phage libraries which display on the surface of the capsid oligopeptides, expressed from random sequence oligonucleotidic inserts introduced into a gene coding for a phage capsid protein using genetic manipulation techniques;

- selection of the phages that display on the surfaces of the capsid antigenic oligopeptides recognized by said antibody; optionally by selection with a first pathologic serum, subsequent screening with a second different pathologic serum and counter-screening with a panel of sera from healthy individuals;

- optional use of the selected phages and/or fragments thereof and/or their derivatives for the formulation of diagnostic kits for the specific pathogenic agent, or in general for the diseases, including immunological disorders typical of so-called autoimmune diseases, with known or unknown etiology and/or pathogenesis;

- optional use of the selected phages and/or fragments thereof and/or their derivatives for the formulation of an antagonist of the antigen-antibody reactions for treatment of the disease induced by said antigen;

- optional use of the selected phages and/or fragments thereof and/or their derivatives to induce a tolerance of the phenomena of hypersensitivity and/or allergy to compounds and/or natural or synthetic preparations;

- 74 -

- optional immunization of an organism by means of the selected phages and/or fragments thereof and/or their derivatives; and

- optional verification of the presence, in the serum of the immunized organism, of antibodies that recognize the above antigen or organism specific to the disease.

2. A process for the preparation of immunogens and diagnostic reagents mimicking an antigen or a pathogenic organism specific to a disease according to claim 1, in which - in the case of said antibodies specific to said pathogenic agent also being protective or neutralizing - the material employed for said immunization is used to formulate a vaccine against said specific pathogenic agent.

3. A process according to claim 1 or 2, in which the antibodies are chosen from the group comprising monoclonal antibodies, polyclonal antibodies and antibodies contained in serum.

4. A process according to claim 3, in which the antibodies are antibodies contained in serum.

5. A process for the preparation of immunogens and diagnostic reagents mimicking an antigen or a pathogenic organism specific to a disease according to claims 1 to 4, in which the construction of phage libraries is performed using filamentous phages chosen from the group comprising M13, F1, Fd, and derivatives thereof.

6. A process for the preparation of immunogens and diagnostic reagents mimicking an antigen or a pathogenic organism specific to a disease according to any one of the preceding claims, in which the gene coding for the phage capsid, with random sequence oligonucleotidic inserts, is the gene coding for the protein VIII of the phage capsid.

7. A process for the preparation of immunogens and

- 75 -

diagnostic reagents mimicking an antigen or a pathogenic organism specific to a disease according to claims 1 to 5, in which the gene coding for the phage capsid, with random sequence oligonucleotidic inserts, is the gene coding for the protein III in the phage capsid.

8. A process for the preparation of immunogens and diagnostic reagents mimicking an antigen or a pathogenic organism specific to a disease according to any one of the preceding claims, in which the pathogenic agent is selected from the group comprising the surface antigen of the virus of human hepatitis B (HBsAg), the virus of human hepatitis C, and antigens pathogenically linked to autoimmune diseases such as diabetes.

9. A process for the preparation of immunogens and diagnostic reagents mimicking an antigen or a pathogenic organism specific to a disease according to claim 8, in which the antibody with neutralizing or protective activity in the face of the pathogenic reagent is contained in the serum of individuals immunized using the surface antigen of the virus of human hepatitis B (HBsAg), or in the serum of individuals infected with the virus of human hepatitis C.

10. A process for the preparation of immunogens and diagnostic reagents mimicking an antigen or a pathogenic organism specific to a disease according to any one of the preceding claims, in which the antigenic oligopeptides recognized by the antibody used are obtained from the expression of random sequence oligonucleotidic inserts, using the plasmid pC89 as a vector.

11. A process for the preparation of immunogens and diagnostic reagents mimicking an antigen or a pathogenic organism specific to a disease according to any one of the preceding claims, in which for the


- 76 -

construction of the phage libraries filamentous phages containing in their capsid, in the protein VIII or in the protein III, from the site identifying the restriction enzyme EcoRI (GAATTC) to that identifying the BamHI restriction enzyme (GGATCC), an amino acid sequence chosen from the group comprising the sequences SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:7, and SEQ ID NO:9 to 68.

12. Phages, characterized by the fact that they show antigenic oligopeptides on the surface of the capsid, and by the fact that they are obtainable during the process for the preparation of immunogens or diagnostic reagents mimicking an antigen or a pathogenic organism specific to a disease according to claims 1 to 11.

13. Vaccines against a specific pathogenic agent, characterized by the fact that they are obtainable using the process according to claims 1 to 11.

14. Modified pC89 plasmids, characterized by the fact that they are usable in the process according to claims 1 to 11 and that they contain, either wholly or in part, a nucleotidic sequence chosen from the group comprising the sequences SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6 and SEQ ID NO:8.



**THIS PAGE BLANK (USPTO)**





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup>:</b> <b>C12N 15/10, 15/62, 15/36, 15/51, 7/01,</b> <b>C07K 7/08, A61K 39/12, G01N 33/577</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 94/26886</b> <b>(43) International Publication Date:</b> 24 November 1994 (24.11.94)
<b>(21) International Application Number:</b> PCT/IT94/00054 <b>(22) International Filing Date:</b> 5 May 1994 (05.05.94) <b>(30) Priority Data:</b> RM93A000301 11 May 1993 (11.05.93) <b>IT</b> <b>(71) Applicant (for all designated States except US):</b> ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P. AN- GELETTI S.P.A. [IT/IT]; Via Pontina Km 30.600, I-00040 Pomezia (IT). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> FELICI, Franco [IT/IT]; Viale Bruno Rizzieri, 247, I-00173 Roma (IT). LUZZAGO, Alessandra [IT/IT]; Via Beata Vergine del Carmelo, 54, I- 00144 Roma (IT). NICOSIA, Alfredo [IT/IT]; Via Beata Vergine del Carmelo, 54, I-00144 Roma (IT). MONACI, Paolo [IT/IT]; Via Appia Nuova, 420, I-00181 Roma (IT). CORTESE, Riccardo [IT/IT]; Via Massimiliano Massimo, 16, I-00144 Roma (IT). <b>(74) Agents:</b> DI CERBO, Mario et al.; Società Italiana Brevetti S.p.A., Piazza di Pietra, 39, I-00186 Roma (IT).	<b>(81) Designated States:</b> AU, BR, CA, CN, JP, RU, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>  <b>(88) Date of publication of the international search report:</b> 16 March 1995 (16.03.95)	

**(54) Title:** PROCESS FOR THE PREPARATION OF IMMUNOGENS OR DIAGNOSTIC REAGENTS, AND IMMUNOGENS OR DIAGNOSTIC REAGENTS THEREBY OBTAINABLE

lac promoter      leader sequence      pVIII      α-peptide

TAG

GAATTC . TAG . GATCCC  
GluPhe . END . AspPro  
EcoRI      BamHI

**(57) Abstract**

A process for the preparation of immunogens or diagnostic reagents that mimic an antigen or a pathogenic organism specific to a disease, essentially characterized by the following operations: identification of at least one antibody that reacts with the antigen or pathogenic organism specific to the disease; construction of phage libraries which display on the surface of the capsid oligopeptides, expressed from random sequence oligonucleotidic inserts introduced into a gene coding for a phage capsid protein using genetic manipulation techniques (for example using a plasmid engineered for the purposes of the invention, the genetic map of which is shown in the figure); selection of the phages that display on the surfaces of the capsid antigenic oligopeptides recognized by said antibody; optional use of the selected phages and/or fragments thereof and/or their derivatives for the formulation of diagnostic kits for the specific pathogenic agent, or in general for the diseases, including immunological disorders typical of so-called autoimmune diseases, with known or unknown etiology and/or pathogenesis; optional use of the selected phages and/or fragments thereof and/or their derivatives for the formulation of an antagonist of the antigen-antibody reactions for treatment of the disease induced by said antigen; optional use of the selected phages and/or fragments thereof and/or their derivatives to induce a tolerance of the phenomena of hypersensitivity and/or allergy to compounds and/or natural or synthetic preparations; optional immunization of an organism by means of the selected phages and/or fragments thereof and/or their derivatives; and optional verification of the presence, in the serum of the immunized organism, of antibodies that recognize the above antigen or organism specific to the disease.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## INTERNATIONAL SEARCH REPORT

 International Application No  
 PCT/IT 94/00054

## A. CLASSIFICATION OF SUBJECT MATTER

 IPC 5 C12N15/10 C12N15/62 C12N15/36 C12N15/51 C12N7/01  
 C07K7/08 A61K39/12 G01N33/577

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C12N C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,91 19818 (AFFYMAX TECHNOLOGIES N.V.) 26 December 1991 see page 6, line 10 - page 28, line 11 see page 7, line 10 - line 19 ---	1-7,10, 12,13
X	TRENDS IN BIOCHEMICAL SCIENCE, vol.17, no.7, July 1992, ELSEVIER SCIENCE, AMSTERDAM, NL; pages 241 - 245 J.K. SCOTT 'Discovering peptide ligands using epitope libraries' see page 245, middle column, line 37 - line 62 ---	1-7,12, 13
X	WO,A,92 07077 (PERHAM, WILLIS) 30 April 1992 Y see page 4, line 34 - page 21, line 13; claims 1-10 ---	1-7,12, 13 10
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*A\* document member of the same patent family

Date of the actual completion of the international search

18 January 1995

Date of mailing of the international search report

1 6. 02 95

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax (+31-70) 340-3016

Authorized officer

Hornig, H

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/IT 94/00054

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J. MOL. BIOL., vol.220, no.4, 1991; ACADEMIC PRESS LIMITED, LONDON, UK; pages 821 - 827	1-7,12, 13
Y	J. GREENWOOD ET AL. 'Multiple display of foreign peptides on a filamentous bacteriophage; Peptides from Plasmodium falciparum Circumsporozoite protein as antigens' cited in the application see page 821, line 1 - line 11 see page 825, left column, line 14 - right column, line 3 ---	10
O,X	GENE, vol.128, 1993, ELSEVIER PUBLISHERS, N.Y., U.S.; pages 85 - 88	1-7,12, 13
Y	O.O. MINENKOVA ET AL. 'Design of specific immunogens using filamentous phages as carrier' Article occasioned by a meeting at the Banbury Center, Cold Spring Harbor Laboratory, NY, USA; April 4-7, 1992; see page 85, left column, line 1 - page 87, right column, line 9 ---	10
O,X	GENE, vol.128, 1993, ELSEVIER PUBLISHERS, N.Y., U.S.; pages 79 - 83	1-7,12, 13
Y	A.E. WILLIS ET AL. 'Immunological properties of foreign peptides in multiple display on a filamentous bacteriophage' Article occasioned by a meeting at the Banbury Center, Cold Spring Harbor Laboratory, NY, USA; April 4-7, 1992; see page 79, left column, line 1 - page 83, left column, line 10 ---	10
O,X	GENE, vol.128, 1993, ELSEVIER PUBLISHERS, N.Y., U.S.; pages 21 - 27	1-7,12, 13
Y	F. FELICI ET AL. 'Mimicking of discontiguous epitopes by phage-displayed peptides, II. Selectin of clones recognized by a protective monoclonal antibody against Bordetella pertussis toxin from phage libraries' Article occasioned by a meeting at the Banbury Center, Cold Spring Harbor Laboratory, NY, USA; April 4-7, 1992 see page 22, right column, line 1 - page 26, left column, line 5 --- -/--	10

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/IT 94/00054

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,93 01288 (DEUTSCHES KREBSFORSCHUNGSZENTRUM STIFTUNG DES ÖFFENTLICHEN RECHTS) 21 January 1993 see page 1, line 10 - page 5, line 21 ---	1-7,10, 12,13
Y	WO,A,92 06176 (IXSYS, INC.) 16 April 1992  see page 5, line 5 - page 27, line 18 ---	1-7,10, 12,13
Y	WO,A,92 06204 (IXSYS, INC.) 16 April 1992  see page 4, line 13 - page 13, line 8 ---	1-7,10, 12,13
Y	WO,A,91 18980 (CETUS CORPORATION) 12 December 1991 see page 1, line 5 - page 12, line 16 ---	1-7,10, 12,13
O,Y	GENE, vol.128, 1993, ELSEVIER PUBLISHERS, N.Y., U.S.; pages 51 - 57 A. LUZZAGO ET AL. 'Mimicking of discontinuous epitopes by phage-displayed peptides, I. Epitope mapping of human H ferritin using a phage library of constrained peptides' cited in the application Article occasioned by a meeting at the Banbury Center, Cold Spring Harbor Laboratory, NY, USA; April 4-7, 1992; see page 51, left column, line 1 - page 56, right column, line 31 ---	1-7,10, 12,13
Y	J. MOL. BIOL., vol.222, 1991, ACADEMIC PRESS LIMITED, LONDON, UK; pages 301 - 310 L. CASTAGNOLI ET AL. 'Selection of antibody ligands from a large library of oligopeptides expressed on a multivalent exposition vector' cited in the application see page 308, right column, line 46 - right column, line 62 --- -/--	1-7,10, 12,13

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	CHEMICAL ABSTRACTS, vol. 121, no. 15, 10 October 1994, Columbus, Ohio, US; abstract no. 176965z, A. FOLGORI ET AL. 'Selection of epitopes from phage displayed peptide libraries by using human sera: a new tool for the identification of antigenic and immunogenic mimotopes' page 830 ;column L ; see abstract & ZENTRALBLATT BAKTERIOL., SUPPL. 1994, 24(BACTERIAL PROTEIN TOXINS),415-425; FEMS Symposium, No.73. Bacterial Protein Toxins; Sixth European Workshop, Stirling Scotland, UK, June 27- July 2, 1993; ---	1-14
T	EMBO JOURNAL, vol.13, no.9, May 1994, IRL PRESS LIM., OXFORD, ENGL.; pages 2236 - 2243 A. FOLGORI ET AL. 'A general strategy to identify mimotopes of pathological antigens using only random peptide libraries and human sera' the whole document ---	1-14
T	TRENDS IN BIOTECHNOLOGY, vol.12, no.7, July 1994, ELSEVIER SCIENCE PUBLISHERS,LTD.,CAMBRIDGE,UK; pages 262 - 267 R. CORTESE ET AL. 'Epitope discovery using peptide libraries displayed on phage' the whole document ---	1-14
T	GENE, vol.146, no.2, 2 September 1994, ELSEVIER PUBLISHERS, N.Y., U.S.; pages 191 - 198 C. MOTTI ET AL. 'Recognition by human sera and immunogenicity of HBsAg mimotopes selected from an M13 phage display library' the whole document -----	1-14

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/IT 94/00054

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see annex

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:  
  
1-10, (11-13) part.  
14
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☒ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/IT 94/ 00054

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

1. Claims: 1-7,10,14 (totally searched)  
8,9,11-13 (partially searched)

Process for the preparation of immunogens or diagnostic reagents that mimic an antigen or a pathogenic organism specific to a disease, essentially characterized by:

- identification of at least one antibody that reacts with the antigen or pathogenic organism specific for the disease;
- construction of a phage libraries which display on the surface of the capsid oligopeptides;
- selection of the phages recognized by said antibody;
- Phages obtainable during said process;
- Vaccines obtainable by said process;

Said process in which the pathogenic organism is the virus hepatitis B; Phages and Vaccines obtained and modified pC89 plasmids obtainable by said process but limited to SEQ ID NO: 4,5,6,8.

2. Claims: 8,9,11-13 (partially searched)

Methods, phages and vaccines as in invention 1, in which the pathogenic agent is selected from hepatitis C (HCV) and/or type II Cryoglobulinemia caused by and/or associated with said virus (SEQ ID NO: 9-42).

3. Claims: 8,11-13 (partially searched)

Methods, phages and products as in invention 1, but limited to the antigen of Type I diabetes autoimmune disease (SEQ ID NO: 43-47).

4. Claims: 11-13 (partially searched)

Methods, phages and products as in invention 1, but limited to the antigen for the detection and induction of the immune response of the human tumoral protein NEU (p185HER2) (SEQ ID NO: 48-68).



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IT 94/00054

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9119818	26-12-91	AU-A- 8285291	07-01-92
		EP-A- 0535151	07-04-93
		JP-T- 5508321	25-11-93
WO-A-9207077	30-04-92	EP-A- 0552267	28-07-93
		JP-T- 6501848	03-03-94
WO-A-9301288	21-01-93	DE-A- 4122599	04-02-93
		EP-A- 0547201	23-06-93
		JP-T- 6500930	27-01-94
WO-A-9206176	16-04-92	AU-A- 8922391	28-04-92
		CA-A- 2092803	29-03-92
		EP-A- 0551438	21-07-93
		JP-T- 6504666	02-06-94
		NZ-A- 239987	26-08-94
WO-A-9206204	16-04-92	AU-A- 8731891	28-04-92
		CA-A- 2092802	29-03-92
		EP-A- 0550645	14-07-93
		JP-T- 6505145	16-06-94
WO-A-9118980	12-12-91	AU-A- 8081491	31-12-91
		EP-A- 0600866	15-06-94

**THIS PAGE BLANK (USPTO)**